

# Trichogin®

(Tinidazole)

500mg TABLETS

ٹرائیکو جین گولیاں

۵۰۰ ملی گرام

TRICHOGIN® Tablets contains Tinidazole a new nitroimidazole derivative distinctively marked by the presence of an ethylsulfonylic chain on the N-ethyl group.

TINIDAZOLE has a remarkably high activity against Trichomonas Vaginalis, Giardia, Amoeba and constitutes a true progress in trichomonicide therapy because it has over other drugs (metronidazole for instance) the following advantages:

- greater trichomonicide activity, both in vitro and in vivo.
- higher and more durable hematic levels.
- excretion through the urinary ways still in active form, which brings in to tact effectively against protozoa dwelling in the urethral duct as well.
- no need for concomitant topical treatment.

TRICHOGIN® is to be administered in a single 2g dose (4 tablets 500 mg taken orally once daily). This mode of administration appears to result in a greater degree to effectiveness than usual treatment schemes, in addition to making the treatment itself simpler, quicker and safer. A single administration leads to quick disappearance of the symptoms and eradication of the pathogenic agent in the vast majority of cases. In the occurrence of trichomonas infection, concomitant treatment of the partner is recommended.

#### INDICATIONS:

Infections from Trichomonas vaginalis, Giardiasis, Amebiasis.

#### DOSAGE:

Trichomoniasis, Giardiasis: 4 tablets as a single dose for once day

Amebiasis 4 tablets as a single dose for 2,3 days.

Administration preferably after meals.

Should the symptoms reappear, administer 4 tablets again in the same way.

#### CAUTION:

For a few days after intake of the tablets, alcoholic drinks should be avoided. No adverse effects of tinidazole on the course of pregnancy have been observed nevertheless TRICHOGIN® like all new drugs, must be used with great caution during pregnancy in view of the fact that some drugs of related nature are apt to cross the placental barrier and get into foetal circulation it seems inadvisable to administer TRICHOGIN® during the first ninety days of pregnancy.

It is recommendable to suspend breast-feeding for a few days after taking TRICHOGIN® since the presence of drugs of similar structure has been detected in the milk of nursing woman as a result of treatment in the course of nursing.

TRICHOGIN® is generally well tolerated occasionally reported disturbances are milk gastro-intestinal upset itching, vertigo. These are always of transient, short duration.

No alteration of laboratory findings have generally been observed as a result of TRICHOGIN® treatment. Metronidazole and other drugs of this group have some times been observed to induce transient leucopenia a similar occurrence might be expected therefore also during treatment with TRICHOGIN®.

#### CONTRAINDICATIONS:

Like all drugs in the nitroimidazole group, TRICHOGIN® is contraindicated in patients with a positive history of blood discrasia, and in patients affected by active organic disease of the central nervous system.

**COMPOSITION:** Each tablet contains Tinidazole B.P. 500 mg.

**PACKING:** Box of 20 tablets in blister packing.

 chiesi

Manufactured by:  
LAHORE CHEMICAL & PHARMACEUTICAL WORKS (PVT.) LTD.  
137-Ferozeshapur Road, Lahore.

Under Licence of  
CHIESI-FARMACEUTICI S.p.A. - ITALY